



### METADATA - IARC Colposcopy Image Bank


Variable name	Variable attributes
CaseID	3 digits
HPV test result	<ul style="list-style-type: none"> <li>• negative</li> <li>• positive</li> <li>• unknown</li> </ul>
Image type	<ul style="list-style-type: none"> <li>• after normal saline</li> <li>• speculum examination</li> <li>• before application of acetic acid</li> <li>• after application of acetic acid</li> <li>• after application of Lugol's iodine</li> </ul>
<b>General assessment:</b>	
Adequacy	<ul style="list-style-type: none"> <li>• adequate</li> <li>• inadequate</li> </ul>
Reason	<ul style="list-style-type: none"> <li>• cervix could not be satisfactorily exposed due to extensive inflammation</li> <li>• stenosis of vagina and atrophy of cervix</li> </ul>
Squamocolumnar junction visibility	<ul style="list-style-type: none"> <li>• completely visible</li> <li>• partially visible</li> <li>• not visible</li> </ul>



Transformation zone type	<ul style="list-style-type: none"> <li>• type 1</li> <li>• type 2</li> <li>• type 3</li> </ul>
<b>Normal colposcopic findings:</b>	
Original squamous epithelium	<ul style="list-style-type: none"> <li>• atrophic</li> <li>• mature</li> </ul>
Columnar epithelium	<ul style="list-style-type: none"> <li>• ectopy absent</li> <li>• ectopy present</li> </ul>
Metaplastic squamous epithelium	<ul style="list-style-type: none"> <li>• crypt gland openings</li> <li>• nabothian cyst</li> <li>• nabothian cyst and crypt gland openings</li> </ul>
Deciduous in pregnancy	<ul style="list-style-type: none"> <li>• no</li> <li>• not applicable</li> </ul>
<b>Abnormal colposcopic findings / General principles / Position and size:</b>	
Location of the lesion	<ul style="list-style-type: none"> <li>• inside the transformation zone (TZ)</li> <li>• outside the transformation zone (TZ)</li> <li>• inside and outside the transformation zone (TZ)</li> <li>• on polyp</li> </ul>
Location of the lesion by clock position	clock position
Additional position	clock position
No. of quadrants of cervix	<ul style="list-style-type: none"> <li>• 1</li> <li>• 2</li> <li>• 3</li> <li>• 4</li> <li>• inside canal</li> <li>• on polyp</li> </ul>



<b>Grade 1 minor:</b>	
Grade 1	<ul style="list-style-type: none"> <li>• fine punctation</li> <li>• irregular border</li> <li>• thin acetowhite epithelium</li> </ul>
Additional Grade 1	<ul style="list-style-type: none"> <li>• fine punctation</li> <li>• irregular border</li> <li>• thin acetowhite epithelium</li> </ul>
<b>Grade 2 major:</b>	
Grade 2	<ul style="list-style-type: none"> <li>• coarse mosaic</li> <li>• coarse punctation</li> <li>• dense acetowhite epithelium</li> <li>• sharp border</li> </ul>
Additional Grade 2	<ul style="list-style-type: none"> <li>• coarse mosaic</li> <li>• coarse punctation</li> <li>• dense acetowhite epithelium</li> <li>• sharp border</li> </ul>
<b>Non-specific:</b>	
Non-specific	<ul style="list-style-type: none"> <li>• erosion</li> <li>• iodine non-stained</li> <li>• iodine stained</li> <li>• leukoplakia</li> </ul>
Additional Non-specific	<ul style="list-style-type: none"> <li>• erosion</li> <li>• iodine non-stained</li> <li>• iodine stained</li> <li>• leukoplakia</li> </ul>
<b>Suspicious for invasion:</b>	
Suspicious for invasion	<ul style="list-style-type: none"> <li>• atypical vessels</li> <li>• irregular surface</li> </ul>
Additional Suspicious for invasion	<ul style="list-style-type: none"> <li>• atypical vessels</li> <li>• irregular surface</li> </ul>




**Miscellaneous finding:**

Miscellaneous finding	<ul style="list-style-type: none"> <li>• condyloma</li> <li>• satellite condyloma</li> <li>• congenital transformation zone</li> <li>• endometriosis</li> <li>• inflammation</li> <li>• ectocervical polyp</li> <li>• endocervical polyp</li> <li>• post-treatment consequence</li> <li>• stenosis</li> </ul>
Additional miscellaneous finding	<ul style="list-style-type: none"> <li>• condyloma</li> <li>• satellite condyloma</li> <li>• congenital transformation zone</li> <li>• endometriosis</li> <li>• inflammation</li> <li>• ectocervical polyp</li> <li>• endocervical polyp</li> <li>• post-treatment consequence</li> <li>• stenosis</li> </ul>

**Swede score:**

Aceto uptake	<ul style="list-style-type: none"> <li>• distinct</li> <li>• stearin/nil or transparent/thin</li> <li>• milky</li> </ul>
Margins	<ul style="list-style-type: none"> <li>• nil or diffuse</li> <li>• sharp but irregular, jagged, satellites</li> <li>• sharp and even, difference in level</li> </ul>
Vessels	<ul style="list-style-type: none"> <li>• absent</li> <li>• coarse or atypical vessels</li> <li>• fine, regular</li> </ul>
Lesion size	<ul style="list-style-type: none"> <li>• &lt; 5 mm/5-15 mm or 2 quadrants</li> <li>• &gt; 15 mm, 3-4 quadrants</li> <li>• endocervically undefined</li> </ul>



Iodine uptake	<ul style="list-style-type: none"><li>• brown</li><li>• distinctly yellow</li><li>• faintly or patchy yellow</li><li>• unknown</li></ul>
Final Swede score	0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 99
<b>Case Summary:</b>	
Provisional diagnosis	diagnosis description
Management	management description
Histopathology	histopathology description